

PHASE TRANSITIONS FOR THE POTTS MODEL WITH FOUR COMPETING INTERACTIONS

HASAN AKIN AND SEYIT TEMIR

ABSTRACT. In this paper we study phase transitions for the Potts model with four competing interactions (external field, the nearest neighbor, the second neighbors and triples of neighbors) on Cayley tree of order two. We prove that for some parameter values of the model there is phase transition. We reduce the problem of describing limiting Gibbs measures to the problem of solving a system of nonlinear functional equations. We extend to the Potts model setting the results obtained by Ganikhodjaev and Rozikov [5] on phase transition for the Ising model.

Mathematics Subject Classification: 82B20, 82B26.

Key words: Potts model; competing interactions; one-level second neighbors.

1. INTRODUCTION

It is well known that lattice spin system is a large class of systems considered in statistical mechanic. Also it is well known that the structure of the lattice plays an important role in examinations of spin systems. The Potts model, which was introduced by Potts [in 1952] as a generalization of the Ising model, has received an increasing theoretical and experimental aspects in recent years (see [13] for details). The idea came from the representation of the Ising model as interactions spin which can be either parallel or anti parallel that is up or down.

We consider a semi-infinite Cayley tree J^k of order $k \geq 2$, i.e., a graph without cycles with $(k+1)$ edges issuing from each vertex except x^0 and with k edges issuing from the vertex x^0 , which is called the tree root. We generalize the problem in [3] for the Potts model with four competing interactions. This problem has been investigated for the Potts model with two competing interactions (see [3] for details).

There are many approaches to derive equation or system equations describing limiting Gibbs measure for the lattice models on Cayley tree (see [1], [10], [11], [12], [13] for details). One of these approaches is based on recursive equations for partition functions ([13]). Another approach is Markov random fields (see [4], [8]). Naturally both approaches lead to the same equation [4]. We concentrate on the first approach, that is, we study recursive equations for partition functions.

In this paper we study the problem of phase transition for the Potts model with four competing interactions (external field, the nearest neighbors, the second neighbors and triples of neighbors) on Cayley tree of order two. In [5], it has been proved that there is phase transition for some parameter values of the Ising model. We generalize this problem for the Potts model. To construct the Hamiltonian equation we consider the generalized Kroneker's symbol δ well coordinated with theory of quadratic stochastic operators.

The rest of the paper is organized as following. In section 2 we give basic definitions and notations. In section 3 we obtain the recursive equations for partition functions. Section 4 contains the solutions of the system of nonlinear equations. The solutions describe whether there is phase transition.

2. PRELIMINARIES

Cayley tree. The Cayley tree Γ^k (see [1]) of order $k \geq 1$ is an infinite tree, i.e., a graph without cycles, from each vertex of which exactly $k + 1$ edges issue. Let $\Gamma^k = (V, L, i)$, where V is the set of vertices of Γ^k , L is the set of edges of Γ^k and i is the incidence function associating each edge $\ell \in L$ with its ends points $x, y \in V$. If $i(\ell) = \{x, y\}$, then x and y are called the nearest neighboring vertices and we write $\ell = \langle x, y \rangle$. For $x, y \in V$, the distance $d(x, y)$ on Cayley tree is defined by formula

$$d(x, y) = \min\{d | x = x_0, x_1, x_2, \dots, x_{d-1}, x_d = y \in V \text{ such that the pairs } \langle x_0, x_1 \rangle, \dots, \langle x_{d-1}, x_d \rangle \text{ are neighboring vertices}\}.$$

For the fixed $x^0 \in V$ we have

$$\begin{aligned} W_n &= \{x \in V : d(x^0, x) = n\} \\ V_n &= \{x \in V : d(x^0, x) \leq n\}, \\ L_n &= \{\ell = \langle x, y \rangle \in L : x, y \in V_n\}, \end{aligned}$$

A collection of the pairs $\langle x_0, x_1 \rangle, \dots, \langle x_{d-1}, x_d \rangle$ is called a path from x to y . We write $x < y$ if the path from x^0 to y goes through x . We call the vertex y a direct successor of x , if $y > x$ and x, y are the nearest neighbors. The set of the direct successors of x is denoted by $S(x)$, i.e.,

$$S(x) = \{y \in W_{n+1} : d(x, y) = 1, x \in W_n\}.$$

It is clear that for any vertex $x \neq x^0$, x has k direct successors and x^0 has $k + 1$. The vertices x and y are called the second neighbor which is denoted by $> x, y <$, if there exists a vertex $z \in V$ such that x, z and y, z are the nearest neighbors.

The second neighbors $> x, y <$ is called one-level neighbors, if vertices x and y belong to W_n for some n , that is, if they are situated on the same level. Three vertices x, y and z are called a triple of neighbors, which is denoted by $\langle x, y, z \rangle$, $\langle x, y \rangle, \langle y, z \rangle$ are nearest neighbors and $x \neq z$. We will consider only one-level the second neighbors.

We assume for neighbors triple $\langle x, y, z \rangle$. For neighbors triple $\langle x, y, z \rangle$ we consider the following Kroneker's symbol

$$(2.1) \quad \delta_{\sigma(x)\sigma(y)\sigma(z)} = \begin{cases} 1, & \text{if } \sigma(x) = \sigma(y) = \sigma(z) \\ \frac{1}{2}, & \text{if } \sigma(x) = \sigma(y) \neq \sigma(z) \text{ or } \sigma(x) \neq \sigma(y) = \sigma(z) \\ 0, & \text{else} \end{cases}$$

where, $x, z \in W_n$ for some n and $y \in W_{n-1}$. We can write (2.1) as follows:

$$\delta_{\sigma(x)\sigma(y)\sigma(z)} = \frac{1}{2} (\delta_{\sigma(x)\sigma(y)} + \delta_{\sigma(y)\sigma(z)}).$$

This definition is well coordinated with theory of quadratic stochastic operators, where quadratic stochastic operator corresponding to generalized Kroneker's symbol in (2.1) is identity transformation [9].

In the Potts model, the spin variables $\sigma(x)$ take their values on a discrete set $\Phi = \{1, 2, \dots, q\}$, $q > 2$ which are associated with each vertex of the tree J^k (see [6] for details).

In this paper, we consider the Potts model with four competing interactions is defined by the following Hamiltonian

$$\begin{aligned}
 H(\sigma) = & -J \sum_{\langle x, y \rangle} \delta_{\sigma(x)\sigma(y)} - J_1 \sum_{>x, y<} \delta_{\sigma(x)\sigma(y)} \\
 (2.2) \quad & - J_2 \sum_{\langle x, y, z \rangle} \delta_{\sigma(x)\sigma(y)\sigma(z)} - h \sum_{x \in V} \delta_{0\sigma(x)}
 \end{aligned}$$

where the sum in the first term ranges all nearest neighbors, second sum ranges all the second neighbors, the third sum ranges all triples of neighbors and the spin variables $\sigma(x)$ assume the values 1, 2, 3. δ is the Kroneker's symbol and J, J_1, J_2 and $h \in \mathbb{R}$ are constant. We assume that $J, J_1, J_2, h \neq 0$. In [6], we solved the problem of the phase transition described by means of the equation (2.2) for $h = 0$ and $J_1 = 0$, it has been found exact solution for the problem of phase transition in the Potts model with competing ternary and binary interactions with external field.

3. RECURSIVE EQUATIONS FOR PARTITION FUNCTIONS

Let Λ be a finite subset of V . We will denote by $\sigma(\Lambda)$ the restriction of σ to Λ . Let $\bar{\sigma}(V \setminus \Lambda)$ be a fixed boundary configuration. The total energy of $\sigma(\Lambda)$ under condition $\bar{\sigma}(V \setminus \Lambda)$ is defined as

$$\begin{aligned}
 H_\Lambda(\sigma(\Lambda) | \bar{\sigma}(V \setminus \Lambda)) = & -J \sum_{<x, y>: x, y \in \Lambda} \delta_{\sigma(x)\sigma(y)} - J \sum_{<x, y>: x \in \Lambda, y \notin \Lambda} \delta_{\sigma(x)\sigma(y)} \\
 & - J_1 \sum_{>x, y<: x, y \in \Lambda} \delta_{\sigma(x)\sigma(y)} - J_1 \sum_{>x, y<: x \in \Lambda, y \notin \Lambda} \delta_{\sigma(x)\sigma(y)} \\
 (3.1) \quad & - J_2 \sum_{<x, y, z>: x, y, z \in \Lambda} \delta_{\sigma(x)\sigma(y)\sigma(z)} - J_2 \sum_{<x, y, z>: x, z \in \Lambda, y \notin \Lambda} \delta_{\sigma(x)\sigma(y)\sigma(z)} \\
 & - h \sum_{x \in \Lambda} \delta_{0\sigma(x)}
 \end{aligned}$$

Then partition function $Z_\Lambda(\bar{\sigma}(V \setminus \Lambda))$ in volume Λ under boundary condition $\bar{\sigma}(V \setminus \Lambda)$ is defined as

$$(3.2) \quad Z_\Lambda(\bar{\sigma}(V \setminus \Lambda)) = \sum_{\sigma(\Lambda) \in \Omega(\Lambda)} \exp(-\beta H_\Lambda(\sigma(\Lambda) | \bar{\sigma}(V \setminus \Lambda))),$$

where $\Omega(\Lambda)$ is the set of all configuration in volume Λ and $\beta = \frac{1}{T}$ is the inverse temperature.

We consider the configurations $\sigma(V_n)$ and the partition functions Z_{V_n} in volume V_n and for the brevity it is denoted them as σ_n and $Z^{(n)}$ respectively ([4], [6]). Let

us decompose the partition function $Z^{(n)}$ into following summands

$$Z^{(n)} = \sum_{i=1}^q Z_i^{(n)},$$

where

$$(3.3) \quad Z_i^{(n)} = \sum_{\sigma_n \in \Omega(V_n): \sigma(x^0)=i} \exp(-\beta H_{V_n}(\sigma_n | \bar{\sigma}(V \setminus V_n))).$$

Let $\theta = \exp(\beta J)$, $\theta_1 = \exp(\beta J_1)$, $\theta_2 = \exp(\beta J_2)$ and $\theta_3 = \exp(\beta h)$.

In this paper, we consider $q = 3$ and $k = 2$. It should be noted that in [3] we have ignored the all triples of neighbors.

Let $S(x^0) = \{x^1, x^2\}$. If $\sigma(x^0) = i$, $\sigma(x^1) = j$ and $\sigma(x^2) = m$, then from (3.1) and (3.2). Thus, we have following equations;

$$Z_i^{(n)} = \sum_{j,m=1}^3 \exp(\beta J(\delta_{ij} + \delta_{im}) + \beta J_1 \delta_{jm} + \beta J_2 \delta_{jim} + h \delta_{1i}) Z_j^{(n-1)} Z_m^{(n-1)}$$

so that

$$\begin{aligned} Z_1^{(n)} &= \theta_3 [\theta^2 \theta_1 \theta_2 (Z_1^{(n-1)})^2 + 2\theta \sqrt{\theta_2} (Z_1^{(n-1)} Z_2^{(n-1)} + Z_1^{(n-1)} Z_3^{(n-1)}) \\ &\quad + \theta_1 ((Z_2^{(n-1)})^2 + (Z_3^{(n-1)})^2) + 2Z_2^{(n-1)} Z_3^{(n-1)}], \end{aligned}$$

$$\begin{aligned} Z_2^{(n)} &= \theta_1 (Z_1^{(n-1)})^2 + 2\theta \sqrt{\theta_2} Z_2^{(n-1)} [Z_1^{(n-1)} + Z_3^{(n-1)}] \\ &\quad + 2Z_1^{(n-1)} Z_3^{(n-1)} + \theta^2 \theta_1 \theta_2 (Z_2^{(n-1)})^2 + \theta_1 (Z_3^{(n-1)})^2 \end{aligned}$$

and

$$\begin{aligned} Z_3^{(n)} &= \theta_1 (Z_1^{(n-1)})^2 + 2\theta \sqrt{\theta_2} Z_3^{(n-1)} [Z_1^{(n-1)} + Z_2^{(n-1)}] \\ &\quad + 2Z_1^{(n-1)} Z_2^{(n-1)} + \theta^2 \theta_1 \theta_2 (Z_3^{(n-1)})^2 + \theta_1 (Z_2^{(n-1)})^2. \end{aligned}$$

After replacement $u_n = \frac{Z_2^{(n)}}{Z_1^{(n)}}$ and $v_n = \frac{Z_3^{(n)}}{Z_1^{(n)}}$ we have the following system of recurrent equations

$$\begin{aligned} \theta_3 u_n &= \frac{\theta_1 + 2\theta \sqrt{\theta_2} u_{n-1} (1 + v_{n-1}) + 2v_{n-1} + \theta^2 \theta_1 \theta_2 u_{n-1}^2 + \theta_1 v_{n-1}^2}{\theta^2 \theta_1 \theta_2 + 2\theta \sqrt{\theta_2} (u_{n-1} + v_{n-1}) + \theta_1 (u_{n-1}^2 + v_{n-1}^2) + 2u_{n-1} v_{n-1}} \\ \theta_3 v_n &= \frac{\theta_1 + 2u_{n-1} + 2\theta \sqrt{\theta_2} v_{n-1} (1 + u_{n-1}) + \theta u_{n-1}^2 + \theta^2 \theta_1 \theta_2 u_{n-1}^2}{\theta^2 \theta_1 \theta_2 + 2\theta \sqrt{\theta_2} (u_{n-1} + v_{n-1}) + \theta_1 (u_{n-1}^2 + v_{n-1}^2) + 2u_{n-1} v_{n-1}} \end{aligned}$$

If $u = \lim u_n$ and $v = \lim v_n$ then we have the following;

$$(3.4) \quad \begin{aligned} \theta_3 u &= \frac{\theta_1 + 2\theta \sqrt{\theta_2} u (1 + v) + 2v + \theta^2 \theta_1 \theta_2 u^2 + \theta_1 v^2}{\theta^2 \theta_1 \theta_2 + 2\theta \sqrt{\theta_2} (u + v) + \theta_1 (u^2 + v^2) + 2uv} \\ \theta_3 v &= \frac{\theta_1 + 2u + 2\theta \sqrt{\theta_2} v (1 + u) + \theta u^2 + \theta^2 \theta_1 \theta_2 u^2}{\theta^2 \theta_1 \theta_2 + 2\theta \sqrt{\theta_2} (u + v) + \theta_1 (u^2 + v^2) + 2uv} \end{aligned}$$

The number of the solutions of equation (3.4) depends on the parameter $\beta = \frac{1}{kT}$. The phase transition usually occurs for low temperature. It is possible to find an

exact value of T^* such that a phase transition occurs for all $T < T^*$ where T^* is called a critical value of temperature.

Finding the exact value of the critical temperature for some models means to exactly solve the models.

4. TRANSLATION-INVARIANT GIBBS MEASURES

In this section we describe the solutions of the system of nonlinear equations

$$(4.1) \quad \begin{aligned} \theta_3 u &= \frac{\theta_1 + 2\theta\sqrt{\theta_2}u(1+v) + 2v + \theta^2\theta_1\theta_2u^2 + \theta_1v^2}{\theta^2\theta_1\theta_2 + 2\theta\sqrt{\theta_2}(u+v) + \theta_1(u^2+v^2) + 2uv} \\ \theta_3 v &= \frac{\theta_1 + 2u + 2\theta\sqrt{\theta_2}v(1+u) + \theta_1u^2 + \theta^2\theta_1\theta_2u^2}{\theta^2\theta_1\theta_2 + 2\theta\sqrt{\theta_2}(u+v) + \theta_1(u^2+v^2) + 2uv} \end{aligned}$$

Note that if there is more than one positive solution for equations (4.1), then there is more than one translation-invariant Gibbs measure corresponding to these solutions. If the equations (4.1) have more than one positive solution, then it is said that a phase transition occurs for the model (3.2) ([5], [6]).

Let us solve the equations (4.1). Subtracting these equations we have

$$(4.2) \quad \theta_3(u-v) = \frac{2(\tilde{\theta}-1)(u-v) + \theta_1(\tilde{\theta}^2-1)(u^2-v^2)}{\tilde{\theta}^2\theta_1 + 2\tilde{\theta}(u+v) + 2uv + \theta_1(u^2+v^2)}$$

so that equality $u = v$ give us some solutions of (4.1), where $\tilde{\theta} = \theta\sqrt{\theta_2}$. For $u = v$ we have

$$(4.3) \quad \theta_3 u = \frac{(\tilde{\theta}^2\theta_1 + 2\tilde{\theta} + \theta_1)u^2 + 2(\tilde{\theta} + 1)u + \theta_1}{2(\theta_1 + 1)u^2 + 4\tilde{\theta}u + \tilde{\theta}^2\theta_1}$$

Let us consider function

$$(4.4) \quad f(u) = \frac{(\tilde{\theta}^2\theta_1 + 2\tilde{\theta} + \theta_1)u^2 + 2(\tilde{\theta} + 1)u + \theta_1}{2(\theta_1 + 1)u^2 + 4\tilde{\theta}u + \tilde{\theta}^2\theta_1}.$$

In order to describe the solution of the system of nonlinear equations in the (4.1), we should analyze the function in the (4.4).

Now we can study elementary analysis of this function as following:

1) Let us take the first derivative of $f(u)$, in this case, if $\tilde{\theta} > 1$, that is $\theta_2 > \frac{1}{\theta^2}$, then $f(u)$ is increasing and if $\tilde{\theta} < 1$, $f(u)$ is decreasing for $u > 0$;

2) Let us take the second derivative of $f(u)$. If $\theta_1 > \theta_1^*$, where

$$\theta_1^* = \frac{\tilde{\theta}^2 + \tilde{\theta} + 1 + \sqrt{9\tilde{\theta}^4 + 26\tilde{\theta}^3 + 35\tilde{\theta}^2 + 50\tilde{\theta} + 33}}{(\tilde{\theta}^2 + 2)(\tilde{\theta} + 1)},$$

then exist inflection point $u^* > 0$ such that $f''(u) > 0$ for $0 < u < u^*$ and $f''(u) < 0$ for $u > u^*$.

Thus we can generalize the Lemma proved in [3] as follows;

Lemma. Let $\tilde{\theta} > 1$, that is $\theta_2 > \frac{1}{\theta^2}$ and there is inflection point. There exist $\eta_1(\theta, \theta_1, \theta_2)$, $\eta_2(\theta, \theta_1, \theta_2)$ with $0 < \eta_1(\theta, \theta_1, \theta_2) < \eta_2(\theta, \theta_1, \theta_2)$ such that the equation (4.1) has three solutions if $\eta_1(\theta, \theta_1, \theta_2) < \theta_3 < \eta_2(\theta, \theta_1, \theta_2)$; has two solutions if

either $\theta_3 = \eta_1(\theta, \theta_1, \theta_2)$ or $\theta_3 = \eta_2(\theta, \theta_1, \theta_2)$ and has single solution in other cases. In fact

$$\eta_i(\theta, \theta_1, \theta_2) = \frac{1}{u_i^*} f(u_i^*),$$

where u_1^*, u_2^* are the solutions of equation

$$uf'(u) = f(u).$$

Proof follows from properties 1) and 2) of function f . Equation $uf'(u) = f(u)$ is equivalent to following equation

$$(4.5) \quad Au^4 + Bu^3 + Cu^2 + Du + E = 0,$$

where $A = 2(\theta_1 + 1)(\tilde{\theta}^2\theta_1 + 2\tilde{\theta} + \theta_1) > 0$, $B = 8(\tilde{\theta} + 1)(\theta_1 + 1) > 0$, $D = 8\tilde{\theta}\theta_1 > 0$, $E = \tilde{\theta}^2\theta_1 > 0$.

As $C = -\tilde{\theta}^4\theta_1^2 - 2\theta_1\tilde{\theta}^3 - \theta_1^2\tilde{\theta}^2 + 8\tilde{\theta}^2 + 8\tilde{\theta} + 6\theta_1^2 + 6\theta_1$, then equation (4.5) has no positive roots if $C > 0$, and have two positive roots u_1^* , u_2^* if $C < 0$. It is easy to show that exist θ_1^{**} such that for all $\theta_1 > \theta_1^{**}$ coefficient C is negative, where $\theta > \sqrt{\frac{2}{\theta_2}}$. Then we have the following Corollary.

Corollary. For $\theta(\theta_2) > \sqrt{\frac{2}{\theta_2}}$, $\theta_1(\theta, \theta_2) > \max(\theta_1^*, \theta_1^{**})$ and $\eta_1(\theta, \theta_1, \theta_2) < \theta_3 < \eta_2(\theta, \theta_1, \theta_2)$ the equation (4.5) has three positive roots. Thus there is phase transition.

Now let us come back to equation (4.2). After cancelling to (u-v) we have

$$(4.6) \quad t = \frac{\theta_1\theta_2s^2 + (2\theta_2\tilde{\theta} - \theta_1(\tilde{\theta}^2 - 1))s + \tilde{\theta}^2\theta_1\theta_2 - 2(\tilde{\theta} - 1)}{2\theta_3(\theta_1 - 1)},$$

where $u + v = s$ and $uv = t$. After dividing the first equation of (4.1) to the second and simplification we have

$$(4.7) \quad t = \frac{\theta_1s^2 + 2s + \theta_1}{\tilde{\theta}^2\theta_1 + \theta_1 - 2\tilde{\theta}},$$

From (4.6) and (4.7) we obtain the following equality;

$$(4.8) \quad \begin{aligned} \frac{\theta_1(\tilde{\theta} + 1)s + 2}{\theta_3} &= \frac{\theta_1[\theta_1(\tilde{\theta} + 1) - 2]s^2 + 2[\theta_1(\tilde{\theta}^2 + \tilde{\theta} - 2) - 2(\tilde{\theta} + 1)]s}{\theta_1(\tilde{\theta}^2 + 1) - 2\tilde{\theta}} \\ &+ \frac{\theta_1[\theta_1(\tilde{\theta}^3 + \tilde{\theta}^2 + 2\tilde{\theta} + 2) - 2(\tilde{\theta}^2 + \tilde{\theta} + 1)]}{\theta_1(\tilde{\theta}^2 + 1) - 2\tilde{\theta}} \end{aligned}$$

It is easy to show that if $\theta_1(\theta, \theta_2) > \frac{2(\tilde{\theta}+1)}{\tilde{\theta}^2+\tilde{\theta}-2}$ then all coefficients in (4.8) are positive.

Note that the system of equations

$$(4.9) \quad \begin{aligned} u + v &= s \\ uv &= t \end{aligned}$$

have solutions when $t < \frac{s^2}{4}$, so that from (4.7) follows

$$(4.10) \quad (\theta_1(\tilde{\theta}^2 - 3) - 2\tilde{\theta})s^2 - 8s - 4\theta_1 > 0$$

Then there is

$$s^* = \frac{4 + \sqrt{16 + 4\theta_1(\theta_1(\tilde{\theta}^2 - 3) - 2\tilde{\theta})}}{\theta_1\tilde{\theta}^2 - 3\theta_1 - 2\tilde{\theta}} > 0,$$

such that for all $s > s^*$ is valid.

Now let us consider equation (4.8). The elementary analysis get following:

1) If

$$\theta_3 < \frac{2[\theta_1(\tilde{\theta}^2 + 1) - 2\tilde{\theta}]}{\theta_1[\theta_1(\tilde{\theta}^3 + \tilde{\theta}^2 + 2\tilde{\theta} + 2) - 2(\tilde{\theta}^2 + \tilde{\theta} + 1)]},$$

then equation (4.8) has single root.

2) If

$$\theta_3 > \frac{2[\theta_1(\tilde{\theta}^2 + 1) - 2\tilde{\theta}]}{\theta_1[\theta_1(\tilde{\theta}^3 + \tilde{\theta}^2 + 2\tilde{\theta} + 2) - 2(\tilde{\theta}^2 + \tilde{\theta} + 1)]}$$

there is k_0 such that the line $y - \frac{2}{\theta_3} = k_0 s$ is a tangent for parabola from right-hand side of (4.8) and then for $\theta_3 < \frac{\theta_1(\tilde{\theta}+1)}{k_0}$ the equation (4.8) has two positive roots and only one of them greater than s^* .

Thus the equation (4.8) has one positive root greater than s^* , when

$$\theta > \sqrt{\frac{3}{\theta_2}},$$

and

$$\frac{2[\theta_1(\tilde{\theta}^2 + 1) - 2\tilde{\theta}]}{\theta_1[\theta_1(\tilde{\theta}^3 + \tilde{\theta}^2 + 2\tilde{\theta} + 2) - 2(\tilde{\theta}^2 + \tilde{\theta} + 1)]} < \theta_3 < \frac{\theta_1(\tilde{\theta} + 1)}{k_0},$$

where k_0 is defined above.

Recalling results for equation (4.3), we have the following:

the system of the equation (4.1) has five solutions if $\theta > \sqrt{\frac{3}{\theta_2}}$, $\theta_1 > A$ and $B < \theta_3 < C$, where

$$(4.11) \quad \begin{aligned} A &= \max(\theta_1^*, \theta_1^{**}) \\ B &= \max(\eta_1, \frac{2[\theta_1(\tilde{\theta}^2 + 1) - 2\tilde{\theta}]}{\theta_1[\theta_1(\tilde{\theta}^3 + \tilde{\theta}^2 + 2\tilde{\theta} + 2) - 2(\tilde{\theta}^2 + \tilde{\theta} + 1)]}) \\ C &= \min(\eta_2, \frac{\theta_1(\tilde{\theta} + 1)}{k_0}). \end{aligned}$$

More detail analysis similar [2] and [3] get that only three solutions are stable.

Thus we have proved the following theorem.

Theorem. Assume the conditions (4.11) are satisfied then for the model (3.1) there are three translation-invariant Gibbs measure, i.e., there is phase transition.

It is easy to show that these three measures correspond to boundary conditions

$$\bar{\sigma}(V \setminus V_n) \equiv i, \quad i = 1, 2, 3.$$

Remark. We can similarly study the solutions of new system of nonlinear equations by using Kroneker's symbol δ that has the form

$$\delta_{\sigma(x)\sigma(y)\sigma(z)} = \begin{cases} 1 & \text{if } \sigma(x) = \sigma(y) = \sigma(z) \\ 0 & \text{else.} \end{cases}$$

REFERENCES

- [1] R. J. Baxter, Exactly solved models in statistical mechanics, London, Academic Press, 1982.
- [2] N. N. Ganikhodjaev, On pure phases of the ferromagnet Potts model with three states on the Bethe lattice of order two, Theor. Math. Phys. **85** (1990) 163-175.
- [3] N. N. Ganikhodjaev, H. Akin, S. Temir, Potts model with two competing binary interactions, Turkish Mathematical Journal (accepted).
- [4] N. N. Ganikhodjaev, C. H. Pah, M. R. Wahiddin, An Ising model with three competing interactions on a Cayley tree, Journal of Math. Phys. **45** (2004) 3645-3658.
- [5] N. N. Ganikhodjaev, U. A. Rozikov, Phase transitions for Ising model four competing interactions, The Abdus salam international centre for theoretical physics, preprint.
- [6] N. N. Ganikhodjaev, S. Temir, H. Akin, The exact solution of the Potts models with external magnetic field on the Cayley tree, CUBO Math. J., **7** (2005) 37-48.
- [7] H. O. Georgii, Gibbs measures and phase transitions, De Gruyter studies in Mathematics, **9**, Berlin, New York, 1988.
- [8] R. Kindermann, J. L. Snell, Markov random fields and their applications, Contemporary Mathematics, **1**, Providence, R. I.: Amer. Math. Soc., 1980.
- [9] A. M. Mariz, C. Tsallis, E. L. Albuquerque, Phase diagram of the Ising model on a Cayley tree in the presence of competing interactions and magnetic fields, Jour. of Stat. Phys., **40** (1985) 577-592.
- [10] K. Preston, Gibbs states on countable sets, London: Cambridge Univ. Press, 1974.
- [11] C. R. da Silca and S. Countiho, Ising model on the Bethe lattice with competing interactions up to the third-nearest-neighbor generation, Phys. Rev. B., **34** (1986) 7975-7985.
- [12] F. Spitzer, Markov random fields on an infinite tree, Ann. Probability, **3** (1975) 387-398.
- [13] F. Y. Wu, The Potts model, Rev. Mod. Phys., **54** (1982) 235-268.

HASAN AKIN, DEPARTMENT OF MATHEMATICS, ARTS AND SCIENCE FACULTY, HARRAN UNIVERSITY, SANLIURFA, 63200, TURKEY

E-mail address: `akinhasan@harran.edu.tr`

SEYIT TEMIR, DEPARTMENT OF MATHEMATICS, ARTS AND SCIENCE FACULTY, HARRAN UNIVERSITY, SANLIURFA, 63200, TURKEY

E-mail address: `temirseyt@harran.edu.tr`